arXiv:1110.2248v1 [math.RA] 11 Oct 2011

Recognizing The Semiprimitivity of $\mathbb{N}$-graded Algebras via Gröbner Bases*

Huishi Li

Department of Applied Mathematics

College of Information Science and Technology

Hainan University

Haikou 570228, China

Abstract. Let $K\langle X\rangle = K\langle X_1,\ldots,X_n\rangle$ be the free K-algebra on $X = \{X_1,\ldots,X_n\}$ over a field K, which is equipped with a weight $\mathbb{N}$-gradation (i.e., each X_i is assigned a positive degree), and let $\mathcal{G}$ be a finite homogeneous Gröbner basis for the ideal $I = \langle\mathcal{G}\rangle$ of $K\langle X\rangle$ with respect to some monomial ordering $\prec$ on $K\langle X\rangle$. It is proved that if the monomial algebra $K\langle X\rangle/\langle\mathbf{LM}(\mathcal{G})\rangle$ is semi-prime, where $\mathbf{LM}(\mathcal{G})$ is the set of leading monomials of $\mathcal{G}$ with respect to $\prec$, then the $\mathbb{N}$-graded algebra $A = K\langle X\rangle/I$ is semiprimitive (in the sense of Jacobson). In the case that $\mathcal{G}$ is a finite non-homogeneous Gröbner basis with respect to a graded monomial ordering $\prec_{gr}$, and the $\mathbb{N}$-filtration FA of the algebra $A = K\langle X\rangle/I$ induced by the $\mathbb{N}$-grading filtration $FK\langle X\rangle$ of $K\langle X\rangle$ is considered, if the monomial algebra $K\langle X\rangle/\langle\mathbf{LM}(\mathcal{G})\rangle$ is semi-prime, then it is proved that the associated $\mathbb{N}$-graded algebra $G(A)$ and the Rees algebra $\widetilde{A}$ of A determined by FA are all semiprimitive.

Key words: Semiprimitive algebra, graded algebra, monomial algebra, Gröbner basis, Ufnarovski graph.

---

*Project supported by the National Natural Science Foundation of China (10971044).

2010 Mathematics Subject Classification: 16W50, 16W70, 16Z05.

1. Introduction

Let $K\langle X\rangle = K\langle X_1,\dots,X_n\rangle$ be the free K-algebra on $X=\{X_1,\dots,X_n\}$ over a field K, which is equipped with a weight $\mathbb{N}$-gradation (i.e., each X_i is assigned a positive degree), and let $\mathcal{G}$ be a finite Gröbner basis for the ideal $I=\langle\mathcal{G}\rangle$ of $K\langle X\rangle$ with respect to some monomial ordering $\prec$ on $K\langle X\rangle$. Consider the algebra $A=K\langle X\rangle/I$, the associated monomial algebra $\overline{A}=K\langle X\rangle/\langle\mathbf{LM}(\mathcal{G})\rangle$ of A (where $\mathbf{LM}(\mathcal{G})$ is the set of leading monomials of $\mathcal{G}$ with respect to $\prec$), the $\mathbb{N}$-filtration FA of A induced by the $\mathbb{N}$-grading filtration of $K\langle X\rangle$, the associated $\mathbb{N}$-graded algebra $G(A)$ and the Rees algebra $\widetilde{A}$ of A determined by FA (see Section 3 for the definitions). In [Li2] and [Li3], it has been proved that many structural properties of $\overline{A}$ can be transferred to A, $G(A)$ and $\widetilde{A}$ (see [Li4] for more details). In this paper, we first show that if $\mathcal{G}$ is a finite homogeneous Gröbner basis and if the monomial algebra $\overline{A}$ is semi-prime, then the $\mathbb{N}$-graded algebra $A=K\langle X\rangle/I$ is semiprimitive (in the sense of Jacobson). In the case that $\mathcal{G}$ is a finite non-homogeneous Gröbner basis with respect to a graded monomial ordering $\prec_{gr}$, if $\overline{A}$ is semi-prime, then we show that the $\mathbb{N}$-graded algebras $G(A)$ and $\widetilde{A}$ are all semiprimitive. Since the semi-primeness of the monomial algebra $\overline{A}$ can be determined in an algorithmic way ([G-IL], [G-I]), our results are algorithmically realizable in case the algorithms given in loc. cit. are implemented on computer.

Throughout this paper, K denotes a field, algebras considered are associative K-algebras with multiplicative identity 1, and ideals considered in an algebra are meant two-sided ideals. For a subset U of an algebra A, we write $\langle U\rangle$ for the ideal generated by U in A. Moreover, we use $\mathbb{N}$, respectively $\mathbb{Z}$, to denote the set of nonnegative integers, respectively the set of integers.

2. Some Known Results on Monomial Algebras

Let $K\langle X\rangle = K\langle X_1,\dots,X_n\rangle$ be the free K-algebra on $X=\{X_1,\dots,X_n\}$, and $\mathcal{B}=\{X_{i_1}^{\alpha_1}X_{i_2}^{\alpha_2}\cdots X_{i_s}^{\alpha_s}\mid X_{i_j}\in X,\ \alpha_j\in\mathbb{N}\}$ the standard K-basis of $K\langle X\rangle$ consisting of all monomials (words) in X_{i_j}'s. For convenience, we use lowercase letters $w,u,v,s,\dots$ to denote monomials in $\mathcal{B}$. In this section we recall from [G-IL] and [G-I] how to recognize the Jacobson semiprimitivity of a finitely presented monomial algebra $R=K\langle X\rangle/\langle\Omega\rangle$ in a computational way, where $\Omega=\{u_1,\dots,u_s\}\subset\mathcal{B}$ is a reduced finite subset of monomials (see the definition below) such that $\Omega\cap X=\emptyset$.

For $u,v\in\mathcal{B}$, we say that v *divides* u, denoted by $v|u$, if $u=wvs$ for some $w,s\in\mathcal{B}$.

We say that a subset $\Omega \subset \mathcal{B}$ is *reduced* if $v, u \in \Omega$ and $v \neq u$ implies $v \not| u$.

2.1. Theorem Let $\Omega = \{u_1, \ldots, u_s\}$ be a reduced finite subset of $\mathcal{B} - X$ and $R = K\langle X\rangle/\langle\Omega\rangle$.
(i) ([G-IL], Theorem 16) The Jacobson radical $J(R)$ of R coincides with the upper nilradical $\mathrm{Nil}(R)$ of R.
(ii) ([G-I], Theorem 2.27) R is semiprimitive (in the sense of Jacobson), i.e., $J(R) = \{0\}$, if and only if R is semi-prime, i.e., $a \in R$ and $aRa = \{0\}$ implies $a = 0$.

□

Let $\Omega = \{u_1, \ldots, u_s\}$ be a reduced finite subset of $\mathcal{B}$, and $\langle\Omega\rangle$ the monomial ideal generated by Ω. Then the set of *normal monomials* (mod $\langle\Omega\rangle$) in $\mathcal{B}$ is defined as

$$N(\Omega) = \{w \in \mathcal{B} \mid u \not| w,\ u \in \Omega\}.$$

For each $u_i \in \Omega$, say $u_i = X_{i_1}^{\alpha_1} \cdots X_{i_r}^{\alpha_r}$ with $X_{i_j} \in X$ and $\alpha_j \in \mathbb{N}$, we write $l(u_i) = \alpha_1 + \cdots + \alpha_r$ for the length of u_i . Put

$$\ell = \max\{\, l(u_i) \mid u_i \in \Omega\}\,.$$

Then the *Ufnarovski graph* of Ω (in the sense of [Uf1]), denoted by $\Gamma(\Omega)$, is defined as a directed graph, in which the set of vertices V is given by

$$V = \{v_i \in N(\Omega) \mid l(v_i) = \ell - 1\}\,,$$

and the set of edges E contains the edge $v_i \to v_j$ if and only if there exist X_k, $X_t \in X$ such that $v_iX_k = X_tv_j \in N(\Omega)$. Since Ω is finite, the directed graph $\Gamma(\Omega)$ is thereby practically constructible.

Remark (i) Note that we have defined the number ℓ above as in [G-IL]. While in [G-I] this number was defined as $m + 1 = \max\left\{ l(u_i) \,\middle|\, u_i \in \Omega\right\}$. So, in the subsequent results we shall use ℓ and $\ell - 1$ instead of $m + 1$ and m.
(ii) To better understand the practical application of $\Gamma(\Omega)$, it is essential to notice that a Ufnarovski graph is defined by using the *length* $l(u)$ *of the monomial* (*word*) $u \in \mathcal{B}$ instead of using the *degree of* u *as a homogeneous element in* $K\langle X\rangle$ whenever a weight $\mathbb{N}$-gradation of $K\langle X\rangle$ is used (see Section 3), though both notions coincide when each X_i is assigned the degree 1.

Basic notions from classical graph theory fully suit an Ufnarofski graph $\Gamma(\Omega)$. For instance, a *route of length* m in $\Gamma(\Omega)$ with $m \geq 1$ is a sequence of edges

$$v_0 \to v_1 \to v_2 \to \cdots \to v_{m-1} \to v_m.$$

If in a route no edge appears repeatedly then it is called a *simple route.* A simple route with $m \geq 1$ and $v_0 = v_m$ is called a *cycle.* If, as an undirected graph, there is a route between any two distinct vertices of $\Gamma(\Omega)$, then $\Gamma(\Omega)$ is called *connected.* A *connected component* of $\Gamma(\Omega)$ (as an undirected graph) is a connected subgraph which is connected to no additional vertices.

Let $\Gamma(\Omega)$ be as above. It follows from [Uf1] that there is a one-to-one correspondence between the monomials (words) of length $\geq \ell - 1$ in $N(\Omega)$ and the routes in $\Gamma(\Omega)$, that is, if $u \in N(\Omega)$ and $u = X_{i_1} \cdots X_{i_t}$ with $t \geq \ell - 1$, then u is mapped to the route

$$\mathscr{R}(u): \quad v_0 \to v_1 \to \cdots \to v_m,$$

where

$$\begin{aligned} &m = t - \ell + 1, \text{ and} \\ &v_j = X_{i_{j+1}} X_{i_{j+2}} \cdots X_{i_{j+\ell-1}},\ 0 \leq j \leq m. \end{aligned}$$

In [G-I] the following notions are introduced. A vertex of $\Gamma(\Omega)$ is called a *cyclic vertex* if it belongs to a cycle. A normal monomial $u \neq 1$ (i.e., $u \in N(\Omega) - \{1\}$) is called a *cyclic normal monomial* if $l(u) \leq \ell - 1$ and u is a suffix of some cyclic vertx of $\Gamma(\Omega)$, or, if $l(u) > \ell - 1$ and the associated route $\mathscr{R}(u)$ of u is a subroute of some cyclic route.

2.2. Proposition Let $\Omega = \{u_1, \ldots, u_s\} \subset \mathcal{B} - X$ be reduced, $R = K\langle X \rangle / \langle \Omega \rangle$, and let $J(R)$ be the Jacobson radical of R. If $u \in N(\Omega) - \{1\}$, then the following statements hold.
(i) ([G-I], Lemma 2.13) u is cyclic if and only if there exists a monomial $v \in \mathcal{B}$ such that $(uv)^q \notin \langle \Omega \rangle$ for all integer $q \geq 0$.
(ii) ([G-I], Corollary 2.17) $\overline{u} \in J(R)$ if and only if u is noncyclic, where $\overline{u}$ is the residue class representaed by u in R.

2.3. Theorem ([G-I], Theorem 2.21) Let the monomial algebra $R = K\langle X \rangle / \langle \Omega \rangle$ be as in Proposition 2.2. Then R is semiprimitive if and only if any monomial $u \in N(\Omega)$ with $1 \leq l(u) \leq \ell$ is cyclic.

□

Remark (i) By Theorem 2.1(ii), it is clear that if $R = K\langle X \rangle / \langle \Omega \rangle$ is a prime ring, then R is semiprimitive.
(ii) The reader is referred to [G-IL] and [G-I] for the algorithms written for determining the semi-primeness and the primeness (and hence the semiprimitivity) of a finitely presented monomial algebra $R = K\langle X \rangle / \langle \Omega \rangle$.

3. The Main Results

In this section, we prove the main results of this paper (Theorem 3.2, Theorem 3.3, Theorem 3.5). The Gröbner basis theory for ideals in a free K-algebra is referred to ([Ber], [Mor], [Gr], [Uf2]).

To begin with, let K be a field and $K\langle X\rangle = K\langle X_1,\dots,X_n\rangle$ the free K-algebra on $X=\{X_1,\dots,X_n\}$. As before the standard K-basis of $K\langle X\rangle$ is denoted by $\mathcal{B}$. We fix a weight $\mathbb{N}$-gradation for $K\langle X\rangle$, that is, $K\langle X\rangle = \oplus_{p\in\mathbb{N}}K\langle X\rangle_p$ in which, each X_i has an assigned positive degree m_i, $1\le i\le n$, and the degree-p homogeneous part $K\langle X\rangle_p$ is the K-vector space spanned by all monomials of degree p. For a nonzero homogeneous element $H\in K\langle X\rangle_p$, we write $d(H)$ for the degree of H, i.e., $d(H)=p$. Note that every monomial $w\in\mathcal{B}$ is a homogeneous element. If I is a graded ideal of $K\langle X\rangle$ (i.e., I is generated by homogeneous elements), then $A=K\langle X\rangle/I$ is an $\mathbb{N}$-graded algebra, that is, $A=\oplus_{p\in\mathbb{N}}A_p$ with the degree-p homogeneous part $A_p=(K\langle X\rangle_p+I)/I$.

Moreover, let $\prec$ be a monomial ordering on $\mathcal{B}$, i.e., $\prec$ is a well-ordering on $\mathcal{B}$ such that $u\prec v$ implies $wus\prec wvs$, and $v=wus$ with $w\ne 1$ or $s\ne 1$ implies $u\prec v$, for all $w,u,v,s\in\mathcal{B}$. With the monomial ordering $\prec$ fixed on $\mathcal{B}$, each subset S of $K\langle X\rangle$ is associated to a subset of monomials $\mathbf{LM}(S)=\{\mathbf{LM}(f)\mid f\in S\}\subset\mathcal{B}$, where if $f\in S$ and $f=\sum_{i=1}^s\lambda_iw_i$ with $\lambda_i\in K$ and $w_i\in\mathcal{B}$ such that $w_1\prec w_2\prec\cdots\prec w_s$, then $\mathbf{LM}(f)=w_s$. $\mathbf{LM}(S)$ is usually referred to as the *set of leading monomials* of S. By the classical Gröbner basis theory of $K\langle X\rangle$, in principle every nonzero ideal I of $K\langle X\rangle$ has a nontrivial (finite or infinite) Gröbner basis $\mathcal{G}$ in the sense that $\mathcal{G}$ is a proper subset of $I-\{0\}$ and $\langle\mathbf{LM}(I)\rangle=\langle\mathbf{LM}(\mathcal{G})\rangle$. A Gröbner basis $\mathcal{G}$ consisting of homogeneous elements of $K\langle X\rangle$ is called a *homogeneous Gröbner basis*.

In proving our main theorems, we need a fundamental result concerning the Jacobson radical of a $\mathbb{Z}$-graded ring, which is due to G. Bergman (cf. [Row]).

3.1. Theorem Let $R=\oplus_{n\in\mathbb{Z}}R_n$ be a $\mathbb{Z}$-graded ring and $J(R)$ the Jacobson radical of R. Then
(i) $J(R)$ is a graded ideal of R; and
(ii) if $n\ne 0$ and $a\in R_n$, then $1+a$ is invertible if and only if a is nilpotent.

□

3.2. Theorem Let $K\langle X\rangle$ and the monomial ordering $\prec$ on $\mathcal{B}$ be as fixed above, and let $\mathcal{G}=\{g_1,\dots,g_s\}$ be a finite homogeneous Gröbner basis for the ideal $I=\langle\mathcal{G}\rangle$, such that $\mathbf{LM}(\mathcal{G})\cap X=\emptyset$ and $\mathbf{LM}(\mathcal{G})$ is reduced (in the sense of Section 2). If the monomial algebra $\overline{A} = K\langle X\rangle/\langle \mathbf{LM}(\mathcal{G})\rangle$ is semi-prime, then the $\mathbb{N}$-graded algebra $A = K\langle X\rangle/I$ is semiprimitive.

Proof Let $N(I)$ be the set of normal monomials (mod I) in $\mathcal{B}$, i.e., $N(I) = \{w \in \mathcal{B} \mid w \notin \langle \mathbf{LM}(I)\rangle\}$. Then, since $\mathcal{G}$ is a Gröbner basis of I, $N(I) = \{u \in \mathcal{B} \mid \mathbf{LM}(g_i) \not| u,\ g_i \in \mathcal{G}\}$. If, with respect to the $\mathbb{N}$-gradation of $K\langle X\rangle$, $H \in K\langle X\rangle_p$ is a homogeneous element of degree p, and if $H \notin I$, then by the division by the homogeneous Gröbner basis $\mathcal{G}$, H has a representation $H = \sum_{i,j} \lambda_{ij} w_{ij} g_j v_{ij} + \sum_t \mu_t u_t$, where $\lambda_{ij}, \mu_t \in K - \{0\}$, $w_{ij}, v_{ij} \in \mathcal{B}$, $g_j \in \mathcal{G}$, and $u_t \in N(I)$ such that $d(H) = p = d(u_t)$ for all t. Putting $H' = \sum_t \mu_t u_t$ and considering the nonzero homogeneous element $\overline{H}$ of degree p represented by H in $A_p = (K\langle X\rangle_p + I)/I$, we have

$$\overline{H} = \overline{H'} = \sum_t \mu_t \overline{u_t}, \quad \mu_t \in K - \{0\},\ u_t \in N(I) \text{ with } d(u_t) = p = d(H). \tag{1}$$

Let $J(A)$ be the Jacobson radical of the $\mathbb{N}$-graded algebra $A = K\langle X\rangle/I$. If $J(A) \neq \{0\}$, then it follows from Theorem 3.1 that $J(A)$ is a graded ideal of A. Taking a nonzero homogeneous element of $J(A)$, say $\overline{H} \in J(A) \cap A_p$, where $A_p = (K\langle X\rangle_p + I)/I$ and $H \in K\langle X\rangle_p$, we may replace $\overline{H}$ by $\overline{H'}$ as in (1) above. Without loss of generality we assume that $\mathbf{LM}(H') = u_1$ with respect to $\prec$. Our aim below is to show that

(•) the normal monomial u_1 is noncyclic, thereby $\overline{u_1} \in J(\overline{A})$ by Proposition 2.2, where $\overline{u_1}$ is the residue class represented by u_1 in $\overline{A}$.

Assume the contrary that u_1 is cyclic (see Section 2 for the definition). Then, by Proposition 2.2, there is a monomial $v \in \mathcal{B}$ such that $(u_1v)^q \notin \langle \mathbf{LM}(I)\rangle = \langle \mathbf{LM}(\mathcal{G})\rangle$, or equivalently, $(u_1v)^q \in N(I)$ for all $q \in \mathbb{N}$. Since $\mathbf{LM}(H'v) = u_1v$, it turns out that

$$\mathbf{LM}\left((H'v)^q\right) = (\mathbf{LM}(H'v))^q = (u_1v)^q \in N(I),\ q \in \mathbb{N}. \tag{2}$$

On the other hand, writing $\overline{v}$ for the residue class represented by v in A, we have $\overline{H'v} = \overline{H'}\,\overline{v} \in J(A)$. As $\overline{H'v}$ is again a homogeneous element of $J(A)$, it follows from Theorem 3.1 that $\overline{H'v}^m = 0$ for some integer $m > 0$. Hence, $(H'v)^m \in I$ and this gives rise to

$$(u_1v)^m = (\mathbf{LM}(H'v))^m = \mathbf{LM}\left((H'v)^m\right) \in \langle \mathbf{LM}(I)\rangle = \langle \mathbf{LM}(\mathcal{G})\rangle. \tag{3}$$

Clearly, (3) contradicts (2). Therefore, u_1 is noncyclic and consequently $\overline{u_1} \in J(\overline{A})$, proving the claim (•).

Finally, suppose that the monomial algebra $\overline{A} = K\langle X\rangle/\langle \mathbf{LM}(\mathcal{G})\rangle$ is semi-prime. Then it follows from Theorem 2.1(ii) that $\overline{A}$ is semiprimitive and hence $J(\overline{A}) = \{0\}$. By the above argument, we conclude that $J(A) = \{0\}$; otherwise, by the claim (•), there would be a normal monomial $u_1 \in N(I)$ such that $\overline{u_1} \in J(\overline{A}) = \{0\}$ and hence, $u_1 \in \langle \mathbf{LM}(\mathcal{G})\rangle$, which is a contradiction. This shows that A is semiprimitive. $\Box$

Let I be an *arbitrary* proper ideal of $K\langle X\rangle$, $A = K\langle X\rangle/I$. Consider the natural $\mathbb{N}$-grading filtration $FK\langle X\rangle = \{F_pK\langle X\rangle\}_{p\in\mathbb{N}}$ of $K\langle X\rangle$ determined by a fixed weight $\mathbb{N}$-gradation for $K\langle X\rangle$, that is, $F_pK\langle X\rangle = \oplus_{q\le p}K\langle X\rangle_q$ for $p \in \mathbb{N}$. Then A has the natural $\mathbb{N}$-filtration $FA = \{F_pA\}_{p\in\mathbb{N}}$ induced by the $\mathbb{N}$-grading filtration $FK\langle X\rangle$, where $F_pA = (F_pK\langle X\rangle + I)/I$ for $p \in \mathbb{N}$, and A has the associated $\mathbb{N}$-graded algebra $G(A) = \oplus_{p\in\mathbb{N}}G(A)_p$ with $G(A)_p = F_pA/F_{<p}A$, where $F_{<p}A = \cup_{q<p}F_qA$ (conventionally we put $F_{<0}A = \{0\}$). In case I is a graded ideal of $K\langle X\rangle$, it is clear that $G(A) \cong A$ as $\mathbb{N}$-graded algebras.

We also recall that any total ordering $\prec$ on $\mathcal{B}$ induces an ordering $\prec_{gr}$ on $\mathcal{B}$ subject to the rule: For $u, v \in \mathcal{B}$,

$$u \prec_{gr} v \Leftrightarrow d(u) < d(v) \text{ or } d(u) = d(v) \text{ and } u \prec v,$$

where $d(\ )$ is the degree function on the homogeneous elements of $K\langle X\rangle$. If $\prec_{gr}$ is a monomial ordering on $\mathcal{B}$, then $\prec_{gr}$ is called an $\mathbb{N}$-*graded monomial ordering*, for instance, the commonly used $\mathbb{N}$-graded lexicographic ordering on $\mathcal{B}$.

3.3. Theorem With notation as before, let $\prec_{gr}$ be an $\mathbb{N}$-graded monomial ordering on $\mathcal{B}$ with respect to a fixed weight $\mathbb{N}$-gradation of $K\langle X\rangle$, and let $\mathcal{G} = \{g_1, \ldots, g_s\}$ be a finite (but not necessarily homogeneous) Gröbner basis for the ideal $I = \langle\mathcal{G}\rangle$, such that $\mathbf{LM}(\mathcal{G}) \cap X = \emptyset$ and $\mathbf{LM}(\mathcal{G})$ is reduced (in the sense of Section 2). Consider the algebra $A = K\langle X\rangle/I$. If the monomial algebra $\overline{A} = K\langle X\rangle/\langle\mathbf{LM}(\mathcal{G})\rangle$ is semi-prime, then $G(A)$ is semiprimitive.

Proof Let $\mathbf{LH}(\mathcal{G}) = \{\mathbf{LH}(g_i) \mid g_i \in \mathcal{G}\}$ be the set of $\mathbb{N}$-leading homogeneous elements of $\mathcal{G}$ with respect to the fixed weight $\mathbb{N}$-gradation of $K\langle X\rangle$, that is, $\mathbf{LH}(g_i) = H_p$ if $g_i = H_0 + H_1 + \cdots + H_p$ with $H_j \in K\langle X\rangle_j$ and $H_p \ne 0$. Since we are using an $\mathbb{N}$-graded monomial ordering $\prec_{gr}$, it follows from ([LWZ], Theorem 2.3.2) or ([Li2], Proposition 3.2) that $\mathbf{LH}(\mathcal{G})$ is a Gröbner basis for the graded ideal $\langle\mathbf{LH}(\mathcal{G})\rangle$ of $K\langle X\rangle$, and that

$$G(A) \cong K\langle X\rangle/\langle\mathbf{LH}(\mathcal{G})\rangle$$

as $\mathbb{N}$-graded algebras. Furthermore, under the $\mathbb{N}$-graded monomial ordering $\prec_{gr}$ we have $\mathbf{LM}(\mathcal{G}) = \mathbf{LM}(\mathbf{LH}(\mathcal{G}))$. Hence the $\mathbb{N}$-graded algebra $K\langle X\rangle/\langle\mathbf{LH}(\mathcal{G})\rangle$ has the associated monomial algebra $\overline{A} = K\langle X\rangle/\langle\mathbf{LM}(\mathcal{G})\rangle$. Consequently, our assertion follows from Theorem 3.2 and the isomorphism given above. $\Box$

Now, we turn to the Rees algebra $\widetilde{A}$ of the $\mathbb{N}$-filtered algebra $A = K\langle X\rangle/I$, where I is an *arbitrary* proper ideal of $K\langle X\rangle$, the $\mathbb{N}$-filtration $FA = \{F_pA\}_{p\in\mathbb{N}}$ for A is as constructed before Corollary 3.3, and $\widetilde{A}$ is defined as the $\mathbb{N}$-graded algebra $\widetilde{A} = \oplus_{p\in\mathbb{N}}F_pA$ with the multiplication induced by $F_pAF_qA \subseteq F_{p+q}A$ for all $p, q \in \mathbb{N}$. The relations between A, $G(A)$ and $\widetilde{A}$ are given by the algebra isomorphisms $A \cong \widetilde{A}/\langle 1 - Z\rangle$ and $G(A) \cong \widetilde{A}/\langle Z\rangle$, where Z is the homogeneous element of degree 1 in $\widetilde{A}_1 = F_1A$ represented by the multiplicative identity element 1 of A. Because of these relations, the structure of $\widetilde{A}$ is closely related to the study of the homogenized algebra of an algebra defined by relations, the regular central extension and the PBW-deformation of an $\mathbb{N}$-graded algebra defined by relations (cf. [LWZ], [Li1], [Li2]).

Consider the free K-algebra $K\langle X, T\rangle = K\langle X_1, \ldots, X_n, T\rangle$ in which each X_i has the same positive degree as fixed in $K\langle X\rangle$, and we assign $d(T) = 1$. Write $\widetilde{\mathcal{B}}$ for the standard K-basis for $K\langle X, T\rangle$. If $\prec_{gr}$ is some $\mathbb{N}$-graded lexicographic ordering on the standard K-basis $\mathcal{B}$ of $K\langle X\rangle$, then $\prec_{gr}$ extends to a $\mathbb{N}$-graded lexicographic ordering $\prec_{_{T\text{-}gr}}$ on $\widetilde{\mathcal{B}}$ subject to $T \prec_{_{T\text{-}gr}} X_i$, $1 \le i \le n$. If $f \in K\langle X\rangle$ has the linear representation $f = \lambda\mathbf{LM}(f) + \sum_i \lambda_i w_i$ with λ, $\lambda_i \in K - \{0\}$, $\mathbf{LM}(f) \in \mathcal{B} \cap K\langle X\rangle_p$, $w_i \in \mathcal{B} \cap K\langle X\rangle_{q_i}$, then the *non-central homogenization* of f with respect to T is the homogeneous element

$$\widetilde{f} = \mathbf{LC}(f)\mathbf{LM}(f) + \sum_i \lambda_i T^{p-q_i} w_i \in K\langle X, T\rangle_p.$$

Clearly, $\mathbf{LM}(f) = \mathbf{LM}(\widetilde{f})$, where $\mathbf{LM}(f)$ is taken with respect to $\prec_{gr}$ on $\mathcal{B}$ and $\mathbf{LM}(\widetilde{f})$ is taken with respect to $\prec_{_{T\text{-}gr}}$ on $\widetilde{\mathcal{B}}$. If I is an ideal of $K\langle X\rangle$, then we put

$$\widetilde{I} = \{\widetilde{f} \mid f \in I\} \cup \{X_iT - TX_i \mid 1 \le i \le n\},$$

and call $\langle\widetilde{I}\rangle$, the graded ideal of $K\langle X, T\rangle$ generated by $\widetilde{I}$, the *non-central homogenization ideal* of I in $K\langle X, T\rangle$ with respect to T.

3.4. Proposition With notation as fixed above, let I be an ideal of $K\langle X\rangle$ and $\mathcal{G} \subset I$. The following statements are equivalent.
(i) $\mathcal{G}$ is a Gröbner basis for I with respect to $\prec_{gr}$ on $\mathcal{B}$.
(ii) $\widetilde{\mathcal{G}} = \{\widetilde{g} \mid g \in \mathcal{G}\} \cup \{X_iT - TX_i \mid 1 \le i \le n\}$ is a homogeneous Gröbner basis for $\langle\widetilde{I}\rangle$ with respect to $\prec_{_{T\text{-}gr}}$ on $\widetilde{\mathcal{B}}$.
(iii) The set of normal monomials (mod$\langle\widetilde{I}\rangle$) in $\widetilde{\mathcal{B}}$, with respect to $\prec_{_{T\text{-}gr}}$, is given by

$$N(\langle\widetilde{I}\rangle) = \{T^r u \mid u \in N(I),\ r \in \mathbb{N}\},$$

where $N(I)$ is the set of normal monomial (mod I) in $\mathcal{B}$ with respect to $\prec_{gr}$.

Proof The equivalence (i) ⇔ (ii) is a strengthened version of ([LWZ], Theorem 2.3.2 (i) ⇔ (ii)), of which a detailed proof was given in [LS].

Noticing that $N(I)=\{u\in\mathcal{B}\mid \mathbf{LM}(g)\not| u,\ g\in\mathcal{G}\}$, $\mathbf{LM}(\mathcal{G})=\mathbf{LM}(\widetilde{\mathcal{G}})$ where $\mathbf{LM}(\mathcal{G})$ is taken with respect to $\prec_{gr}$ on $\mathcal{B}$ and $\mathbf{LM}(\widetilde{\mathcal{G}})$ is taken with respect to $\prec_{_{T\text{-}gr}}$ on $\widetilde{\mathcal{B}}$, and that $\mathbf{LM}(X_iT-TX_i)=X_iT$ with respect to $\prec_{_{T\text{-}gr}}$ on $\widetilde{\mathcal{B}}$, the verification of the equivalence (ii) $\Leftrightarrow$ (iii) is straightforward by referring to the well-known characterization of a Gröbner basis in terms of the remainder on division by $\widetilde{\mathcal{G}}$. $\Box$

With the preparation made above, we are ready to mention and prove the next

3.5. Theorem With notation as fixed above, let $\mathcal{G}=\{g_1,\ldots,g_s\}$ be a finite (but not necessarily homogeneous) Gröbner basis for the ideal $I=\langle\mathcal{G}\rangle$ with respect to $\prec_{gr}$ on $\mathcal{B}$, such that $\mathbf{LM}(\mathcal{G})\cap X=\emptyset$ and $\mathbf{LM}(\mathcal{G})$ is reduced (in the sense of Section 2). Consider the algebra $A=K\langle X\rangle/I$ which has the $\mathbb{N}$-filtration FA as constructed before. If the monomial algebra $\overline{A}=K\langle X\rangle/\langle\mathbf{LM}(\mathcal{G})\rangle$ is semi-prime, then the Rees algebra $\widetilde{A}$ of A is semiprimitive.

Proof First note that if the $\mathbb{N}$-graded lexicographic ordering $\prec_{gr}$ on $\mathcal{B}$ is defined subject to

$$X_{i_1}\prec_{gr}X_{i_2}\prec_{gr}\cdots\prec_{gr}X_{i_n},$$

then the $\mathbb{N}$-graded lexicographic ordering $\prec_{_{T\text{-}gr}}$ on $\widetilde{\mathcal{B}}$ is defined subject to

$$T\prec_{_{T\text{-}gr}}X_{i_1}\prec_{_{T\text{-}gr}}X_{i_2}\prec_{_{T\text{-}gr}}\cdots\prec_{_{T\text{-}gr}}X_{i_n}.$$

Moreover, we also bear in mind that $d(T)=1$, and that each X_i has the same positive degree as fixed in $K\langle X\rangle$.

Let $\langle\widetilde{I}\rangle$ be the non-central homogenization ideal of I in $K\langle X,T\rangle$ with respect to T. Then, by Proposition 3.4(iii), the set of normal monomials (mod$\langle\widetilde{I}\rangle$) in $\widetilde{\mathcal{B}}$ with respect to $\prec_{_{T\text{-}gr}}$ is given by $N(\langle\widetilde{I}\rangle)=\{T^ru\mid u\in N(I),\ r\in\mathbb{N}\}$, where $N(I)$ is the set of normal monomial (mod I) in $\mathcal{B}$ with respect to $\prec_{gr}$. If $T^ru_1,\ T^su_2\in N(\langle\widetilde{I}\rangle)$ with $d(T^ru_1)=d(T^su_2)$ such that $T^ru_1\prec_{_{T\text{-}gr}}T^su_2$, then it follows from the definition of $\prec_{_{T\text{-}gr}}$ that

$$r\geq s \text{ and } \ u_1\prec_{gr}u_2. \tag{1}$$

Consider the $\mathbb{N}$-gradation of $K\langle X,T\rangle$ determined by the assigned degrees for T and X_i's. If $H\in K\langle X,T\rangle_p$ is a homogeneous element of degree p, and if $H\not\in\langle\widetilde{I}\rangle$, then since $\widetilde{\mathcal{G}}=\{\widetilde{g}\mid g\in\mathcal{G}\}\cup\{X_iT-TX_i\mid 1\leq i\leq n\}$ is a homogeneous Gröbner basis for $\langle\widetilde{I}\rangle$ with respect to $\prec_{_{T\text{-}gr}}$ (Proposition 3.4(ii)), the division by $\widetilde{\mathcal{G}}$ yields a representation $H=D+\sum_i\lambda_iT^{r_i}u_i$, where $D\in\langle\widetilde{I}\rangle$, $\lambda_i\in K-\{0\}$, and $T^{r_i}u_i\in N(\widetilde{I})$ such that $d(T^{r_i}u_i)=p=d(H)$ for all i. Put $H'=\sum_i\lambda_iT^{r_i}u_i$ and consider the $\mathbb{N}$-graded algebra

$K\langle X,T\rangle/\langle\widetilde{I}\rangle=\oplus_{p\in\mathbb{N}}(K\langle X,T\rangle_p+\langle\widetilde{I}\rangle)/\langle\widetilde{I}\rangle$. Then the nonzero homogeneous element $\overline{H}$ of degree p represented by H in $(K\langle X,T\rangle_p+\langle\widetilde{I}\rangle)/\langle\widetilde{I}\rangle$ has the representation

$$\overline{H}=\overline{H'}=\sum_i\lambda_i\overline{T^{r_i}u_i},\quad \lambda_i\in K-\{0\},\ u_i\in N(I)\text{ with }d(T^{r_i}u_i)=p=d(H).\tag{2}$$

Moreover, By the above (1), if $\mathbf{LM}(H')=T^{r_1}u_1$ with respect to $\prec_{_{T\text{-}gr}}$, then

$$\mathbf{LM}\left(\sum_i\lambda_iu_i\right)=u_1\text{ with respect to }\prec_{gr}.\tag{3}$$

By the definition of $\widetilde{I}$, it is not difficult to verify that the map

$$\begin{array}{rccccl}\psi: & K\langle X,T\rangle/\langle\widetilde{I}\rangle & \longrightarrow & K\langle X\rangle/I=A & \\ & \overline{T} & \mapsto & 1 & \\ & \overline{X_i} & \mapsto & \overline{X_i} & 1\le i\le n\end{array}$$

is well defined, and that ψ is a K-algebra epimorphism. If, as described in (2) above, $\overline{H}=\overline{H'}=\sum_i\lambda_i\overline{T^{r_i}u_i}$ is a nonzero homogeneous element of degree p in $K\langle X,T\rangle/\langle\widetilde{I}\rangle$, then since $u_i\in N(I)$ and conventionally the ideal I considered is a proper ideal, we have

$$\psi(\overline{H})=\psi(\overline{H'})=\sum_i\lambda_i\overline{u_i}=\overline{\sum_i\lambda_iu_i}\neq 0.\tag{4}$$

By ([LWZ], [Li1], [Li2]), $\widetilde{A}\cong K\langle X,T\rangle/\langle\widetilde{I}\rangle$ as $\mathbb{N}$-graded algebras, that is, the algebra isomorphism gives rise to isomorphisms of K-vector spaces

$$\widetilde{A}_p\cong(K\langle X,T\rangle_p+\langle\widetilde{I}\rangle)/\langle\widetilde{I}\rangle,\quad p\in\mathbb{N}.$$

Identifying $\widetilde{A}$ with $K\langle X,T\rangle/\langle\widetilde{I}\rangle$, we now proceed to deal with the Jacobson radical $J(\widetilde{A})$ of $\widetilde{A}$. If $J(\widetilde{A})\neq\{0\}$, then $J(\widetilde{A})$ is a graded ideal of $\widetilde{A}$ by Theorem 3.1. Taking a nonzero homogeneous element of $J(\widetilde{A})$, say $\overline{H}\in J(\widetilde{A})\cap\widetilde{A}_p$, where $\widetilde{A}_p=(K\langle X,T\rangle_p+\langle\widetilde{I}\rangle)/\langle\widetilde{I}\rangle$ and $H\in K\langle X,T\rangle_p$, we may replace $\overline{H}$ by $\overline{H'}$ as in (2) above. Without loss of generality we assume that $\mathbf{LM}(H')=T^{r_1}u_1$. Our aim below is to show that

(∗) the normal monomial u_1 is noncyclic, thereby $\overline{u_1}\in J(\overline{A})$ by Proposition 2.2, where $\overline{u_1}$ is the residue class represented by u_1 in $\overline{A}$.

Assume the contrary that u_1 is cyclic (see Section 2 for the definition). Then, by Proposition 2.2, there is a monomial $v\in\mathcal{B}$ such that $(u_1v)^q\notin\langle\mathbf{LM}(\mathcal{G})\rangle$ for all $q\in\mathbb{N}$, or equivalently,

$$(u_1v)^q\in N(I)\text{ for all }q\in\mathbb{N},\text{ in particular }u_1v\in N(I).\tag{5}$$

On the other hand, $\overline{H'v} = \overline{H'}\overline{v} \in J(\widetilde{A})$. As $\overline{H'v}$ is again a homogeneous element of $J(A)$, it follows from Theorem 3.1 that

$$\overline{H'v}^m = 0 \text{ for some integer } m > 0, \text{ i.e., } (H'v)^m \in \langle \widetilde{I} \rangle. \tag{6}$$

Furthermore, put $H'' = \sum_i \lambda_i u_i v$. By the foregoing (3), $\mathbf{LM}(H'') = u_1 v$ with respect to $\prec_{gr}$. Since $\psi(\overline{H'}) \neq 0$ and $u_1 v \in N(I)$ by (4) and (5), we have $\psi(\overline{H'v}) = \overline{H''} \neq 0$ in A. But it follows from (6) that $\overline{H''}$ is a nilpotent element in A, i.e.,

$$\overline{H''}^m = \left(\sum_i \lambda_i \overline{u_i v} \right)^m = \left(\psi(\overline{H'v}) \right)^m = \psi\left(\overline{H'v}^m \right) = 0. \tag{7}$$

Hence, $(H'')^m \in I$ and this gives rise to

$$(u_1 v)^m = (\mathbf{LM}(H''))^m = \mathbf{LM}((H'')^m) \in \langle \mathbf{LM}(I) \rangle = \langle \mathbf{LM}(\mathcal{G}) \rangle. \tag{8}$$

Clearly, (8) contradicts (5). Therefore, u_1 is noncyclic and consequently $\overline{u_1} \in J(\overline{A})$, proving the claim $(*)$.

Finally, suppose that the monomial algebra $\overline{A} = K\langle X \rangle / \langle \mathbf{LM}(\mathcal{G}) \rangle$ is semi-prime. Then it follows from Theorem 2.1(ii) that $\overline{A}$ is semiprimitive and hence $J(\overline{A}) = \{0\}$. By the above argument, we conclude that $J(\widetilde{A}) = \{0\}$; otherwise, by the claim $(*)$, there would be a normal monomial $u_i \in N(I)$ such that $\overline{u_1} \in J(\overline{A}) = \{0\}$ and hence, $u_1 \in \langle \mathbf{LM}(\mathcal{G}) \rangle$, which is a contradiction. This shows that $\widetilde{A}$ is semiprimitive. □

We end this paper by the following

Open question Let the K-algebra $A = K\langle X \rangle / \langle \mathcal{G} \rangle$ be as in Theorem 3.5. If the monomial algebra $\overline{A} = K\langle X \rangle / \langle \mathbf{LM}(\mathcal{G}) \rangle$ is semi-prime, is A semiprimitive?

References

[Ber3] G. Bergman, The diamond lemma for ring theory, *Adv. Math.*, 29(1978), 178–218.

[G-I] T. Gateva-Ivanova, Algorithmic determination of the Jacobson radical of monomial algebras, in: *Proc. EUROCAL'85*, LNCS Vol. 378, Springer-Verlag, 1989, 355–364.

[G-IL] T. Gateva-Ivanova and V. Latyshev, On recognizable properties of associative algebras, *J. Symbolic Computation*, 6(1988), 371–388.

[Gr] E. Green, An introduction to noncommutative Gröbner bases, in: *Computational Algebra*, Proceedings of the fifth meeting of the Mid-Atlantic Algebra Conference, 1993, (K. G. Fischer, P. Loustaunau, J. Shapiro, E. L. Green, and D. Farkas eds.), Lecture Notes in Pure and Applied Mathematics, Vol. 151, Marcel Dekker, 1994, 167–190.

[Li1] H. Li, *Noncommutative Gröbner Bases and Filtered-Graded Transfer*, Lecture Notes in Mathematics, Vol. 1795, Springer, 2002.

[Li2] H. Li, Γ-leading homogeneous algebras and Gröbner bases, in: *Recent Developments in Algebra and Related Areas* (F. Li and C. Dong eds.), Advanced Lectures in Mathematics, Vol. 8, International Press & Higher Education Press, Boston-Beijing, 2009, 155 – 200. arXiv:math.RA/0609583, http://arXiv.org

[Li3] H. Li, On the calculation of gl.dim$G^{\mathbb{N}}(A)$ and gl.dim$\widetilde{A}$ by using Gröbner bases, *Algebra Colloquium*, 16(2)(2009), 181–194. arXiv:math.RA/0805.0686, http://arXiv.org

[Li4] H. Li, *Gröbner Bases in Ring Theory*, Monograph, World Scientific Publishing Co., Oct. 2011.

[LS] H. Li and C. Su, On (de)homogenized Gröbner bases, *Journal of Algebra, Number Theory: Advances and Applications*, 3(1)(2010), 35–70.
arXiv:0907.0526, http://arXiv.org

[LWZ] H. Li, Y. Wu and J. Zhang, Two applications of noncommutative Gröbner bases, *Ann. Univ. Ferrara - Sez. VII - Sc. Mat.*, XLV(1999), 1–24.

[Mor] T. Mora, An introduction to commutative and noncommutative Gröbner Bases, *Theoretic Computer Science*, 134(1994), 131–173.

[Row] L.H. Rowen, *Ring Theory*, Vol. I, Pure and Applied Mathematics vol. 127, Academic Press, 1988.

[Uf1] V. Ufnarovski, On the use of graphs for computing a basis, growth and Hilbert series of associative algebras, (in Russian 1989), *Math. USSR Sbornik*, 180(11)(1989), 417-428.

[Uf2] V. Ufnarovski, Introduction to noncommutative Gröbner basis theory, in: *Gröbner Bases and Applications* (Linz, 1998), London Math. Soc. Lecture Notes Ser., 251, Cambridge Univ. Press, Cambridge, 1998, 259–280.